

STATE DOCUMENTS COLLECTION

MAR 23 1996

MONTANA STATE LIBRARY  
1515 E. 6TH AVE.  
HELENA, MONTANA 59620

**Aquatic Macroinvertebrate Rapid Bioassessment  
of Soda Butte Creek, Park County, MT.  
July 21, 1994.**

Prepared for the  
Montana Department of Health & Environmental Sciences  
Water Quality Division

Prepared by  
Daniel L. McGuire  
Aquatic Biologist  
Espanola, NM

February 6, 1995

Due

MAY 01 1996

MAY 12 1996

## Introduction

Concern for water quality and aquatic resources in Soda Butte Creek has prompted the Montana Department of Health and Environmental Sciences - Water Quality Division (WQD) to continue environmental studies of this stream. In 1994, environmental monitoring included water chemistry, periphyton and aquatic macroinvertebrates at five locations in the Soda Butte Creek drainage.

This report presents and analyzes the macroinvertebrate data. The objectives of this report are to (1) provide a current assessment of biotic condition at five stream sites (2) describe the degree and probable causes of biological impairment at each site and (3) develop a baseline for future monitoring.

## Rationale

Aquatic macroinvertebrate communities consist primarily of immature insects, including stoneflies (Plecoptera), caddisflies (Trichoptera), mayflies (Ephemeroptera), true flies (Diptera), beetles (Coleoptera) and others. These organisms are important components of aquatic ecosystems, and are the energy links between primary producers (algae), organic inputs to the stream and fish. Macroinvertebrates are good indicators of environmental conditions due to their limited mobility, predictable associations with specific habitats, and differential tolerances to pollution. Evaluating the biological integrity of this assemblage can provide an assessment of environmental quality and can be used to identify limiting factors, for detecting impacts from physical alterations, sediment deposition, nutrients and toxicants, and to document successful mitigation of environmental degradation.

Macroinvertebrate-based Rapid Bioassessment Protocols (RBP) surveys provide a general analysis of biological integrity. This multiple-metric approach quantifies attributes of community composition, structural and functional organization into a single number estimate of biological health. Biological integrity is defined as "the capability of supporting and maintaining a balanced, integrated, adaptive community having species composition, diversity and functional organization comparable to that of natural habitat of the region" (Karr and Dudley 1981).



## Study Area

Aquatic macroinvertebrates were collected at four locations on Soda Butte Creek and at one location on Republic Creek in Park County, Montana. Station designations are:

- SB-1 Soda Butte Creek above Cook City
- SB-2 Soda Butte Creek below Cook City
- SB-3 Soda Butte Creek above Silver Gate
- SB-4 Soda Butte Creek below Silver Gate
- Re-1 Republic Creek at confluence with Soda Butte Creek

Republic Creek flows into Soda Butte Creek just below station SB-2.

## Methods

Field work was conducted by WQD personnel on 21 July, 1994. At each site, a single collection of macroinvertebrates was obtained using standard traveling kick-net methods (Bukantis 1994). Sampling was conducted after runoff at flows ranging from approximately 10 cfs at the upper site to approximately 20 cfs at the downstream site. Riparian vegetation was generally lacking in the study area.

Laboratory and data analyses were contracted to McGuire Consulting. Techniques described in the most recent Montana RBP guidelines (Bukantis 1994) were followed. RBP III sorting methodology (Plafkin et al. 1989) was employed to obtain a 300 organism subsample from each kick-net collection. Macroinvertebrates were identified to the lowest practical taxonomic level, usually genus or species. Tolerance values and functional designations (Appendix A) used in metric calculations were those recommended by Bukantis (1994).



Macroinvertebrate data were analyzed to provide a numerical estimate of biological integrity. For RBP analyses, Montana streams have been grouped into three primary physiographic regions: mountains, foothills and valleys, and plains (Bahls et al. 1992). Metrics and scoring criteria have been developed for each ecoregion (McGuire 1994). The Soda Butte Creek samples were evaluated using the mountain ecoregion criteria which incorporates seven metrics (Table 1). Each metric received a score ranging from 0 (severely impaired) to 6 (nonimpaired). Scores for all metrics were totaled and biological integrity was expressed as a percentage of the maximum possible score. Impairment classifications (Table 2) were from Plafkin et al. (1989).

A variety of other metrics were also calculated (Appendix B), including a metals tolerance index (McGuire 1993). While not incorporated into the RBP assessment, these metrics were used to further describe the community and supplement baseline data.

**Table 1. Metrics and scoring criteria for 300 organism RBP kick samples from mountain streams in Montana.**

| Mountain metric      | Scoring Criteria |       |       |     |
|----------------------|------------------|-------|-------|-----|
|                      | 6                | 4     | 2     | 0   |
| Taxa richness        | >28              | 28-24 | 24-19 | <19 |
| EPT richness         | >19              | 19-18 | 17-16 | <16 |
| Biotic index         | <3               | 3 - 4 | 4 - 5 | >5  |
| % dominant taxon     | <25              | 25-35 | 35-45 | >45 |
| % Collector-FFG      | <60              | 60-70 | 70-80 | >80 |
| % Scrapers+Shredders | >55              | 55-40 | 40-25 | <25 |
| % EPT                | >70              | 70-55 | 55-40 | <40 |

**Table 2. Criteria for the assessment of biologically significant environmental degradation (from Plafkin et al 1989).**

| % comp. to reference | Classification      |
|----------------------|---------------------|
| >83%                 | nonimpaired         |
| 54-79%               | slightly impaired   |
| 21-50%               | moderately impaired |
| <17%                 | severely impaired   |





## Results and Discussion

Based on mountain ecoregion criteria, RBP assessments indicated slight biological impairment at all five sites (Table 3). Biointegrity estimates for Soda Butte Creek ranged from 81% above Cook City (SB-1) to 57% below Cook City (SB-2). Biological integrity improved gradually downstream from Cook City. Biointegrity was 62 and 67%, respectively, above (SB-3) and below (SB-4) Silver Gate. Biointegrity in Republic Creek, which flows into Soda Butte Creek below SB-2, was 57%.

Symptoms of biological impairment were consistent throughout the study area. At each site, biointegrity was limited by the simplicity of the community and impairment was indicated by structural metrics: taxa richness, EPT richness and the percent relative abundance of the dominant taxa. In contrast, metrics quantifying community composition and functional organization generally received maximum scores.

Habitat degradation and low productivity appeared to be primary causes of biological impairment. During July 1994, the truncated structure and low diversity of the macroinvertebrate community may have been due to armored substrates, low substrate heterogeneity and limited food resources. The low relative abundance of macroinvertebrates that feed on plant debris (shredders) indicated minimal organic input from riparian zones.

The species inhabiting Soda Butte Creek were indicative of excellent water quality and metrics based on indicator-organisms received maximum scores. The metals tolerance index and the biotic index (sensitive to organic pollution) were both extremely low (Appendix B). Benthic community composition appeared typical of a soft-water mountain stream with low productivity. Heptageniid mayflies were numerically dominant at each station.



Among all sites, SB-1 had the highest biointegrity. The benthic community was more diverse and taxa richness was substantially higher at this site than at other sites in the drainage. SB-1 was located immediately below reclaimed mine tailings. It is possible that iron bacteria growing in the stream augmented macro-invertebrate food sources at this site. Both the metals tolerance index and the biotic index were substantially higher at SB-1 than at downstream sites; however, neither metric indicated significant pollution or water quality degradation.

Since Soda Butte Creek has a history of metals pollution, the results of this investigation are somewhat surprising. The macro-invertebrate taxa in the study area were considered highly sensitive to metals pollution and the weight of evidence suggested that metals pollution was not a serious problem during 1994. However, a few metrics could be interpreted as indicating slight metals-related impacts. Low EPT richness is often considered an indicator of metals pollution (McGuire 1993). EPT richness was depressed in both Soda Butte and Republic creeks. In addition, low macroinvertebrate densities can also be indicative of metals pollution and the semi-quantitative (timed) kick samples collected during this study indicated low community density at SB-2 compared to the other sites. Thus, additional macroinvertebrate and water quality investigations may be warranted.



**Table 3. Metric values and bioassessments for Soda Butte and Republic creeks, Park County, MT. July 21, 1994.**

| metric                 | s t r e a m             |              |              |              |              |
|------------------------|-------------------------|--------------|--------------|--------------|--------------|
|                        | Republic                | Soda Butte 1 | Soda Butte 2 | Soda Butte 3 | Soda Butte 4 |
| Taxa richness          | 18                      | 31           | 15           | 15           | 20           |
| EPT richness           | 13                      | 16           | 10           | 14           | 17           |
| Biotic index           | 0.6                     | 2.0          | 0.3          | 0.6          | 0.5          |
| % dominant taxon       | 47                      | 30           | 61           | 42           | 49           |
| % Collector-FFG        | 19                      | 40           | 8            | 24           | 17           |
| % Scrapers + Shredders | 71                      | 46           | 80           | 74           | 78           |
| % EPT                  | 96                      | 78           | 94           | 99           | 93           |
| metric                 | m e t r i c   s c o r e |              |              |              |              |
|                        | Republic                | Soda Butte 1 | Soda Butte 2 | Soda Butte 3 | Soda Butte 4 |
| Taxa richness          | 0                       | 6            | 0            | 0            | 2            |
| EPT richness           | 0                       | 2            | 0            | 0            | 2            |
| Biotic index           | 6                       | 6            | 6            | 6            | 6            |
| % dominant taxon       | 0                       | 4            | 0            | 2            | 0            |
| % Collector-FFG        | 6                       | 6            | 6            | 6            | 6            |
| % Scrapers + Shredders | 6                       | 4            | 6            | 6            | 6            |
| % EPT                  | 6                       | 6            | 6            | 6            | 6            |
| total score            | 24                      | 34           | 24           | 26           | 28           |
| % of reference         | 57%                     | 81%          | 57%          | 62%          | 67%          |
| classification         | SLI                     | SLI          | SLI          | SLI          | SLI          |

\* classifications: (NON) nonimpaired, (SLI) slightly impaired, (MOD) moderately impaired,



## LITERATURE CITED

- Bahls, L., R. Bukantis and S. Tralles. 1992. Benchmark biology of Montana reference streams. Montana Dept. of Health & Environmental Sciences.
- Barbour, M.T., J.L. Plafkin, B.P. Bradley, C.G. Graves and R.W. Wisseman. 1992. Evaluation of EPA's Rapid Bioassessment benthic metrics: metric redundancy and variability among reference streams. *Environmental Toxicology and Chemistry*. 11:437-449.
- Bukantis, R. 1994. DRAFT - Rapid bioassessment macroinvertebrate protocols: sampling and sample analysis SOP's. (May, 1994). Montana Dept. of Health & Environmental Sciences.
- McGuire, D. L. 1993. Clark Fork River macroinvertebrate community biointegrity, 1986 through 1992. Final Report to Montana Dept. of Health and Environmental Sciences.
- McGuire, D. L. 1994. Montana reference streams: 1993 aquatic macroinvertebrate surveys. Prepared for the Montana Dept. of Health & Environmental Sciences - Water Quality Bureau.





Appendix A. Checklist of aquatic macroinvertebrates from Soda Butte Creek, Park County, MT. 21 July 94 with tolerance values for the biotic index, metals tolerance index and functional feeding group designations.

| Taxon                        | BI | MTI | FFG          |
|------------------------------|----|-----|--------------|
| <b>COLEOPTERA</b>            |    |     |              |
| Dytiscidae                   |    |     |              |
| <i>Deronectes sp.</i>        | 5  | 7   | predator     |
| <b>DIPTERA</b>               |    |     |              |
| Chironomidae                 |    |     |              |
| <i>Diamesa sp.</i>           | 5  | 9   | Col-gatherer |
| <i>Pagastia sp.</i>          | 1  | 9   | Col-gatherer |
| <i>Cricotopus sp.</i>        | 7  | 9   | Col-gatherer |
| <i>Orthocladius sp.</i>      | 6  | 5   | Col-gatherer |
| <i>Paraphaenocladus sp.</i>  | 4  | 4   | Col-gatherer |
| <i>Rheocricotopus sp.</i>    | 4  | 5   | Col-gatherer |
| <i>Tvetenia sp.</i>          | 5  | 4   | Col-gatherer |
| <i>Micropsectra sp.</i>      | 4  | 1   | Col-gatherer |
| <i>Tanytarsus sp.</i>        | 6  | 3   | Filterer     |
| Tipulidae                    |    |     |              |
| <i>Hexatoma sp.</i>          | 2  | 2   | predator     |
| <i>Rhabdomastix sp.</i>      | 1  | 1   | Col-gatherer |
| Ceratopogoninae              | 6  | 5   | predator     |
| Simuliidae                   |    |     |              |
| <i>Prosimulium sp.</i>       | 4  | 2   | Filterer     |
| <b>EPHEMEROPTERA</b>         |    |     |              |
| Baetidae                     |    |     |              |
| <i>Baetis bicaudatus</i>     | 2  | 4   | Col-gatherer |
| Ephemerellidae               |    |     |              |
| <i>Drunella coloradensis</i> | 0  | 0   | Scraper      |
| <i>Drunella doddsi</i>       | 1  | 0   | Scraper      |
| <i>Serratella sp.</i>        | 2  | 1   | Col-gatherer |
| Heptageniidae                |    |     |              |
| <i>Cinygmula sp.</i>         | 0  | 0   | Scraper      |
| <i>Epeoris deceptivus</i>    | 0  | 0   | Scraper      |
| <i>Epeorus longimanus</i>    | 1  | 0   | Scraper      |
| <i>Rhithrogena sp.</i>       | 0  | 2   | Scraper      |
| Siphonuridae                 |    |     |              |
| <i>Ameletus sp.</i>          | 0  | 1   | Col-gatherer |



Appendix A. Checklist of aquatic macroinvertebrates from Soda Butte Creek, Park County, MT. 21 July 94 with tolerance values for the biotic index, metals tolerance index and functional feeding group designations.

| Taxon                            | BI | MTI | FFG          |
|----------------------------------|----|-----|--------------|
| PLECOPTERA                       |    |     |              |
| Nemouridae                       |    |     |              |
| <i>Visoka cataractae</i>         | 0  | 0   | Shredder     |
| <i>Zapada oregonensis</i> gp     | 3  | 3   | Shredder     |
| Perlodidae                       |    |     |              |
| <i>Isoperla</i> sp.              | 2  | 3   | predator     |
| <i>Kogotus</i> sp.               | 1  | 2   | predator     |
| <i>Megarcys</i> sp.              | 1  | 1   | predator     |
| Chloroperlinae                   | 2  | 1   | predator     |
| Perlidae                         |    |     |              |
| <i>Doroneuria</i> sp.            | 0  | 2   | predator     |
| TRICHOPTERA                      |    |     |              |
| Hydropsychidae                   |    |     |              |
| <i>Parapsyche</i> sp.            | 0  | 1   | Filterer     |
| Limnephilidae                    |    |     |              |
| <i>Ecclisomyia</i> sp.           | 4  | 2   | Col-gatherer |
| <i>Oligophlebodes</i> sp.        | 3  | 1   | Scraper      |
| Glossosomatidae                  |    |     |              |
| <i>Glossosoma</i> sp.            | 0  | 2   | Scraper      |
| Rhyacophilidae                   |    |     |              |
| <i>Rhyacophila alberta</i> gp.   | 0  | 1   | predator     |
| <i>Rhyacophila brunnea</i> gp.   | 2  | 1   | predator     |
| <i>Rhyacophila hyalinata</i> gp. | 0  | 1   | predator     |
| <i>Rhyacophila verrula</i> gp.   | 0  | 1   | predator     |
| <i>Rhyacophila vofixa</i> gp.    | 0  | 1   | predator     |
| ANNELIDA                         |    |     |              |
| Enchytraeidae                    | 4  | 1   | Col-gatherer |
| Turbellaria                      | 4  | 4   | predator     |



## AQUATIC MACROINVERTEBRATE DATA

Soda Butte &amp; Republic creeks, RBP kick samples 21 Jul 94

|                                | Stream: Republic |    | Soda Butte |     |     | MEAN  | Ave. Dev. | Relative %Diff |
|--------------------------------|------------------|----|------------|-----|-----|-------|-----------|----------------|
| Taxon                          | at mouth         | 1  | 2*         | 3   | 4   |       |           |                |
| COLEOPTERA                     |                  |    |            |     |     |       |           |                |
| <i>Deronectes sp.</i>          | 0                | 1  | 0          | 0   | 0   | 0.2   | 0.3       | 160%           |
| DIPTERA                        |                  |    |            |     |     |       |           |                |
| <i>Diamesa sp.</i>             | 0                | 3  | 0          | 0   | 0   | 0.6   | 1.0       | 160%           |
| <i>Pagastia sp.</i>            | 0                | 2  | 0          | 0   | 0   | 0.4   | 0.6       | 160%           |
| <i>Cricotopus sp.</i>          | 1                | 21 | 0          | 0   | 0   | 4.4   | 6.6       | 151%           |
| <i>Orthocladius sp.</i>        | 0                | 2  | 0          | 0   | 0   | 0.4   | 0.6       | 160%           |
| <i>Paraphaenocladius sp.</i>   | 0                | 1  | 0          | 0   | 0   | 0.2   | 0.3       | 160%           |
| <i>Rheocricotopus sp.</i>      | 1                | 5  | 1          | 0   | 0   | 1.4   | 1.4       | 103%           |
| <i>Tvetenia sp.</i>            | 3                | 1  | 1          | 0   | 0   | 1.0   | 0.8       | 80%            |
| <i>Micropsectra sp.</i>        | 5                | 2  | 5          | 2   | 18  | 6.4   | 4.6       | 73%            |
| <i>Tanytarsus sp.</i>          | 0                | 14 | 0          | 0   | 0   | 2.8   | 4.5       | 160%           |
| <i>Hexatoma sp.</i>            | 0                | 0  | 0          | 0   | 1   | 0.2   | 0.3       | 160%           |
| <i>Rhabdomastix sp.</i>        | 0                | 1  | 1          | 0   | 0   | 0.4   | 0.5       | 120%           |
| Ceratopogoninae                | 0                | 3  | 0          | 0   | 0   | 0.6   | 1.0       | 160%           |
| <i>Prosimulium sp.</i>         | 0                | 1  | 0          | 0   | 0   | 0.2   | 0.3       | 160%           |
| HEMEROPTERA                    |                  |    |            |     |     |       |           |                |
| <i>Baetis bicaudatus</i>       | 38               | 47 | 6          | 71  | 27  | 37.8  | 17.0      | 45%            |
| <i>Drunella coloradensis</i>   | 9                | 8  | 5          | 66  | 17  | 21.0  | 18.0      | 86%            |
| <i>Drunella doddsi</i>         | 0                | 0  | 0          | 5   | 2   | 1.4   | 1.7       | 120%           |
| <i>Serratella sp.</i>          | 0                | 0  | 0          | 0   | 1   | 0.2   | 0.3       | 160%           |
| <i>Cinygmula sp.</i>           | 128              | 85 | 134        | 128 | 145 | 124.0 | 15.6      | 13%            |
| <i>Epeorus deceptivus</i>      | 36               | 2  | 36         | 19  | 54  | 29.4  | 15.1      | 51%            |
| <i>Epeorus longimanus</i>      | 0                | 0  | 0          | 1   | 0   | 0.2   | 0.3       | 160%           |
| <i>Rhithrogena sp.</i>         | 13               | 0  | 0          | 2   | 4   | 3.8   | 3.8       | 99%            |
| <i>Ameletus sp.</i>            | 0                | 5  | 1          | 0   | 0   | 1.2   | 1.5       | 127%           |
| PLECOPTERA                     |                  |    |            |     |     |       |           |                |
| <i>Visoka cat.</i>             | 0                | 3  | 0          | 0   | 0   | 0.6   | 1.0       | 160%           |
| <i>Zapada oregonensis gp</i>   | 5                | 33 | 0          | 1   | 5   | 8.8   | 9.7       | 110%           |
| <i>Isoperla sp.</i>            | 2                | 1  | 0          | 0   | 0   | 0.6   | 0.7       | 120%           |
| <i>Kogotus sp.</i>             | 0                | 1  | 0          | 0   | 1   | 0.4   | 0.5       | 120%           |
| <i>Megarcys sp.</i>            | 5                | 0  | 1          | 1   | 1   | 1.6   | 1.4       | 85%            |
| Chloroperlinae                 | 0                | 7  | 2          | 0   | 0   | 1.8   | 2.2       | 120%           |
| <i>Doroneuria sp.</i>          | 0                | 1  | 0          | 0   | 0   | 0.2   | 0.3       | 160%           |
| TRICHOPTERA                    |                  |    |            |     |     |       |           |                |
| <i>Parapsyche sp.</i>          | 5                | 6  | 3          | 0   | 3   | 3.4   | 1.7       | 49%            |
| <i>Agallomyia sp.</i>          | 0                | 1  | 0          | 0   | 0   | 0.2   | 0.3       | 160%           |
| <i>Oligophlebodes sp.</i>      | 2                | 0  | 0          | 2   | 2   | 1.2   | 1.0       | 80%            |
| <i>Glossosoma sp.</i>          | 0                | 0  | 0          | 2   | 1   | 0.6   | 0.7       | 120%           |
| <i>Rhyacophila alberta gp.</i> | 3                | 6  | 1          | 0   | 3   | 2.6   | 1.7       | 65%            |
| <i>Rhyacophila brunnea gp.</i> | 0                | 0  | 0          | 1   | 0   | 0.2   | 0.3       | 160%           |



# AQUATIC MACROINVERTEBRATE DATA

Soda Butte & Republic creeks, RBP kick samples 21 Jul 94

| Taxon                            | Stream: Republic |    | Soda Butte |   |   | MEAN | Ave. Dev. | Relative %Diff |
|----------------------------------|------------------|----|------------|---|---|------|-----------|----------------|
|                                  | at mouth         | 1  | 2*         | 3 | 4 |      |           |                |
| <i>Rhyacophila hyalinata</i> gp. | 14               | 2  | 0          | 1 | 1 | 3.6  | 4.2       | 116%           |
| <i>Rhyacophila verrula</i> gp.   | 0                | 0  | 0          | 0 | 1 | 0.2  | 0.3       | 160%           |
| <i>Rhyacophila vofixa</i> gp.    | 2                | 15 | 17         | 3 | 5 | 8.4  | 6.1       | 72%            |
| ANNELIDA                         |                  |    |            |   |   |      |           |                |
| Enchytraeidae                    | 0                | 1  | 0          | 0 | 0 | 0.2  | 0.3       | 160%           |
| OTHERS                           |                  |    |            |   |   |      |           |                |
| Turbellaria                      | 1                | 5  | 5          | 0 | 1 | 2.4  | 2.1       | 87%            |

## METRICS

|                            |      |      |      |      |      |      |      |      |
|----------------------------|------|------|------|------|------|------|------|------|
| TOTAL ORGANISMS            | 273  | 286  | 219  | 305  | 293  | 275  | 23.4 | 8%   |
| TAXA RICHNESS              | 18   | 31   | 15   | 15   | 20   | 19.8 | 4.6  | 23%  |
| EPT RICHNESS               | 13   | 16   | 10   | 14   | 17   | 14.0 | 2.0  | 14%  |
| % EPT                      | 96   | 78   | 94   | 99   | 93   | 92   | 6    | 6%   |
| BIOTIC INDEX               | 0.57 | 1.98 | 0.31 | 0.55 | 0.54 | 0.8  | 0.5  | 60%  |
| % DOMINANT TAXON           | 47   | 30   | 61   | 42   | 49   | 46   | 8.0  | 17%  |
| % CHIRONOMIDAE             | 4    | 18   | 3    | 1    | 6    | 6    | 4.6  | 73%  |
| Baetidae-Ephemeropter      | 17   | 32   | 3    | 24   | 11   | 17   | 8.5  | 49%  |
| % COLLECTORS (g+f)         | 19   | 40   | 8    | 24   | 17   | 22   | 8.1  | 38%  |
| % SCRAPER+SHREDDER         | 71   | 46   | 80   | 74   | 78   | 70   | 9.6  | 14%  |
| SHANNON DIVERSITY          | 2.70 | 3.58 | 2.02 | 2.23 | 2.50 | 3    | 0.4  | 16%  |
| METALS TOLERANCE INDEX     | 0.97 | 2.47 | 0.38 | 1.00 | 0.60 | 1    | 0.6  | 51%  |
| EPT/(EPT + Chironomidae)   | 0.96 | 0.81 | 0.97 | 0.99 | 0.94 | 1    | 0.0  | 5%   |
| % FILTERERS                | 2    | 7    | 1    | 0    | 1    | 2    | 2.0  | 87%  |
| % COLLECTOR-GATHERERS      | 18   | 32   | 7    | 24   | 16   | 19   | 7.0  | 37%  |
| % SHREDDERS                | 2    | 13   | 0    | 0    | 2    | 3    | 3.7  | 113% |
| % SCRAPERS                 | 69   | 33   | 80   | 74   | 77   | 67   | 13.3 | 20%  |
| % PREDATORS                | 10   | 15   | 12   | 2    | 5    | 9    | 4.2  | 49%  |
| Scraper/(Scraper+Filterer) | 0.97 | 0.82 | 0.98 | 1.00 | 0.99 | 0.95 | 0.05 | 6%   |
|                            |      |      |      |      |      |      |      |      |
| % COLEOPTERA               | 0    | 0    | 0    | 0    | 0    | 0    | 0.1  | 160% |
| % DIPTERA                  | 4    | 20   | 4    | 1    | 6    | 7    | 5.1  | 75%  |
| % EPHEMEROPTERA            | 82   | 51   | 83   | 96   | 85   | 80   | 11.2 | 14%  |
| % PLECOPTERA               | 4    | 16   | 1    | 1    | 2    | 5    | 4.4  | 89%  |
| % TRICHOPTERA              | 10   | 10   | 10   | 3    | 5    | 8    | 2.7  | 36%  |
| % MISC                     | 0    | 2    | 2    | 0    | 0    | 1    | 0.9  | 92%  |
|                            |      |      |      |      |      |      |      |      |
| # of scraper taxa          | 5    | 3    | 3    | 8    | 7    | 5    | 1.8  | 35%  |
| # of shredder taxa         | 1    | 2    | 0    | 1    | 1    | 1    | 0.4  | 40%  |
| # of predator taxa         | 6    | 10   | 5    | 4    | 8    | 7    | 1.9  | 29%  |
| # of C-gatherer taxa       | 5    | 13   | 6    | 2    | 3    | 6    | 3.0  | 51%  |

\* entire sample used





AQUATIC MACROINVERTEBRATE DATA RBP KICK SAMPLES  
Soda Butte & Republic creeks, RBP kick samples 21 Jul 94

% Relative abundance

|                   | Republic | SB1 | SB2 | SB3 | SB4 |
|-------------------|----------|-----|-----|-----|-----|
| <b>COLEOPTERA</b> |          |     |     |     |     |
| Deronectes sp.    | 0%       | 0%  | 0%  | 0%  | 0%  |

**DIPTERA**

|                      |    |    |    |    |    |
|----------------------|----|----|----|----|----|
| Diamesa sp.          | 0% | 1% | 0% | 0% | 0% |
| Pagastia sp.         | 0% | 1% | 0% | 0% | 0% |
| Cricotopus sp.       | 0% | 7% | 0% | 0% | 0% |
| Orthocladus sp.      | 0% | 1% | 0% | 0% | 0% |
| Paraphaenocladus sp. | 0% | 0% | 0% | 0% | 0% |
| Rheocricotopus sp.   | 0% | 2% | 0% | 0% | 0% |
| Tvetenia sp.         | 1% | 0% | 0% | 0% | 0% |
| Microspectra sp.     | 2% | 1% | 2% | 1% | 6% |
| Tanytarsus sp.       | 0% | 5% | 0% | 0% | 0% |
| Hexatoma sp.         | 0% | 0% | 0% | 0% | 0% |
| Rhabdomastix sp.     | 0% | 0% | 0% | 0% | 0% |
| Ceratopogoninae      | 0% | 1% | 0% | 0% | 0% |
| Prosimulium sp.      | 0% | 0% | 0% | 0% | 0% |

**HEMEROPTERA**

|                       |     |     |     |     |     |
|-----------------------|-----|-----|-----|-----|-----|
| baetis bicaudatus     | 14% | 16% | 3%  | 23% | 9%  |
| Drunella coloradensis | 3%  | 3%  | 2%  | 22% | 6%  |
| Drunella doddsi       | 0%  | 0%  | 0%  | 2%  | 1%  |
| Serratella sp.        | 0%  | 0%  | 0%  | 0%  | 0%  |
| Cinygmula sp.         | 47% | 30% | 61% | 42% | 49% |
| Epeoris deceptivus    | 13% | 1%  | 16% | 6%  | 18% |
| Epeoris longimanus    | 0%  | 0%  | 0%  | 0%  | 0%  |
| Rhithrogena sp.       | 5%  | 0%  | 0%  | 1%  | 1%  |
| Ameletus sp.          | 0%  | 2%  | 0%  | 0%  | 0%  |

**PLECOPTERA**

|                       |    |     |    |    |    |
|-----------------------|----|-----|----|----|----|
| Visoka cat.           | 0% | 1%  | 0% | 0% | 0% |
| Zapada oregonensis gp | 2% | 12% | 0% | 0% | 2% |
| Isoperla sp.          | 1% | 0%  | 0% | 0% | 0% |
| Kogotus sp.           | 0% | 0%  | 0% | 0% | 0% |
| Megarcys sp.          | 2% | 0%  | 0% | 0% | 0% |
| Chloroperlinae        | 0% | 2%  | 1% | 0% | 0% |
| Doroneuria sp.        | 0% | 0%  | 0% | 0% | 0% |

**TRICHOPTERA**

|                         |    |    |    |    |    |
|-------------------------|----|----|----|----|----|
| Parapsyche sp.          | 2% | 2% | 1% | 0% | 1% |
| Trichopsyche sp.        | 0% | 0% | 0% | 0% | 0% |
| Trichophlebodes sp.     | 1% | 0% | 0% | 1% | 1% |
| Glossosoma sp.          | 0% | 0% | 0% | 1% | 0% |
| Rhyacophila alberta gp. | 1% | 2% | 0% | 0% | 1% |
| Rhyacophila brunnea gp  | 0% | 0% | 0% | 0% | 0% |



AQUATIC MACROINVERTEBRATE DATA RBP KICK SAMPLES  
Coda Butte & Republic creeks, RBP kick samples 21 Jul 94

% Relative abundance

|                         | Republic | SB1 | SB2 | SB3 | SB4 |
|-------------------------|----------|-----|-----|-----|-----|
| Rhyacophila hyalinata g | 5%       | 1%  | 0%  | 0%  | 0%  |
| Rhyacophila verrula gp. | 0%       | 0%  | 0%  | 0%  | 0%  |
| Rhyacophila vofixa gp.  | 1%       | 5%  | 8%  | 1%  | 2%  |
| Enchytraeidae           | 0%       | 0%  | 0%  | 0%  | 0%  |
| OTHERS                  |          |     |     |     |     |
| Turbellaria             | 0%       | 2%  | 2%  | 0%  | 0%  |

| stations | QSI-taxa | DIC-5 | QSI-FFG |
|----------|----------|-------|---------|
| 1 vs 2   | 56%      | 2     | 65%     |
| 1 vs 3   | 71%      | 3     | 87%     |
| 1 vs 4   | 69%      | 3     | 77%     |
| 1 vs 5   | 82%      | 3     | 94%     |
| 2 vs 3   | 47%      | 3     | 53%     |
| 2 vs 4   | 52%      | 2     | 66%     |
| 2 vs 5   | 50%      | 2     | 60%     |
| 3 vs 4   | 55%      | 4     | 65%     |
| 3 vs 5   | 77%      | 4     | 87%     |
| 4 vs 5   | 68%      | 4     | 77%     |
| min      | 47%      | 2     | 53%     |
| max      | 82%      | 4     | 94%     |
| mean     | 63%      | 3     | 73%     |

